

广东黄杨科一新种

叶华谷

(中国科学院华南植物研究所, 广东 广州 510650)

A New Species of Buxaceae from Guangdong

YE Hua-gu

(South China Institute of Botany, the Chinese Academy of Sciences, Guangzhou 510650, China)

关键词: 黄杨科; 黄杨属; 广东

Key words: Buxaceae; *Buxus*; Guangdong

中图分类号: Q949.754.1 文献标识码: A 文章编号: 1005-3395(2002)03-0245-02

潮安黄杨 新种 图 1

Buxus chaoanensis H. G. Ye, sp. nov. Fig. 1

Species floribus femineis inter stylos 3 glandibus praeditis insignis, *B. cephalantha* Lev. et Vant. var. *shantouensi* M. Cheng affinis, sed foliis obovato-spathulatis, crasse coriaceis, nervis lateralibus utrinque obsoletis differt. Type: Guangdong, Chaoan Xian, in crevices of rocks of valleys, alt. 300 m. H. G. Ye 319 (holo-, IBSC).

Evergreen shrub, 10–30(–40) cm high. Branchlet angulate, glabrous. Leaves opposite; petiole 1–2 mm, papillose; leaf blade thickly leathery, obovate to narrowly obovate, 1–1.5 cm long and (3–) 4–7 mm broad, apex rounded, emarginate, base cuneate, decurrent into a short petiole, margin entire, ±revolute; midrib raised on both sides; lateral nerves inconspicuous. Inflorescence terminal and axillary, capitate; rhachis 4–6 mm long, puberulent, with 1–2 pairs of bract at base; bracts narrowly ovate-triangular, 1.5 mm × 0.6 mm, apex acuminate. Male flower 5–6 in each inflorescence, subsessile; Sepal 4, triangular-ovate, cymbiform, 1.5–2 mm × 0.8 mm; anther c. 1 mm; filament flat; reduced pistil small, style c. 2 mm. Female flower solitary at apex of the inflorescence; petal 6, ovate, cymbiform, c. 2 mm × 1 mm, apex acute, ciliate; ovary ovoid, c. 2.5 mm, with 3 large glands among the styles; style 3 erect, c. 1 mm; stigma obovate, 2 lobed. Capsule ellipsoid, 5–6 mm long and 4.5 mm broad, apex emarginate. Flowering: April; fruiting: Aug.–Oct.

Specimens examined: Guangdong, Chaoan Xian. In crevices of rocks in valleys, alt: 300 m.
F. Y. Zeng 1516.

收稿日期: 2001-12-06 接受日期: 2002-05-08

基金项目: 广东省自然科学基金资助项目(970848).

常绿灌木，高 10–30(–40)cm。小枝具棱，无毛。叶对生，叶片倒卵形或狭倒卵形，革质，长 1–1.5 cm，宽(3–)4–7 mm，顶端圆钝，微凹，基部楔形，下延成短柄，两面无毛，边缘全缘，反卷，中脉在两面凸起，侧脉不明显；叶柄长 1–2 mm，两侧具乳头状突起。头状花序腋生兼顶生，花序轴长 4–6 mm，被微柔毛；花序轴基部苞片 1–2 对，狭卵状三角形，长约 1.5 mm，宽约 0.6 mm，顶端渐尖；雄花：6–8 朵，近无梗，萼片 4 枚，成 2 对，三角状卵形，凹陷呈舟状，长 1.5–2 mm，宽约 0.8 mm，花药长约 1 mm，花丝扁平，退化子房小，花柱长约 2 mm；雌花 1 朵，位于花序轴顶端，萼片 6 枚，卵形，凹陷呈舟状，长约 2 mm，宽约 1.5 mm，顶端急尖，边缘具缘毛，子房卵球形，长约 2.5 mm，顶端中央具 3 枚明显的圆形大腺体，花柱 3 枚，直立，长约 1 mm，柱头狭倒心形，2 深裂。蒴果椭圆形，长 5–6 mm，宽约 4.5 mm，顶端凹陷；宿存腺体顶端凹陷。花期：4 月；果期：8–10 月。

本种与汕头黄杨 (*B. cephalantha* var. *shantouensis* M. Cheng) 相近，主要区别在于花柱与子房顶端之间具 3 个大腺体，叶革质，匙状倒卵形。

A distinctive species, in many ways similar to *B. cephalantha* Lévl. et Vant. var. *shantouensis* M. Cheng, differs mainly in having three large glands among styles at the top of the ovary, and thickly leathery obovate-spathulate leaves.




图 1 Fig. 1 潮安黄杨 *Buxus chaoanensis* H. G. Ye

1. 花枝 Flowering branches; 2. 花序 Inflorescence; 3. 雄花 Male flower; 4. 雌花 Female flower; 5. 果实 Fruit;
6, 7. 叶片上面 Adaxial surface of leaf blade; 8, 9. 叶片背面 Abaxial surface of leaf blade (刘运笑绘 Drawn by LIU Yun-xiao)